

EXERCISE AS MEDICINE  
WOEMA WEBINAR 12/16/10  
SELECTED REFERENCES

American Thoracic Society. ATS Statement; Guidelines for the six-minute walk test. *Am J Respir Crit Care Med.* 2002;166;111-7.

Arena R; Myers J; Guazzi M. The future of aerobic exercise testing in clinical practice: Is it the ultimate vital sign? *Future Cardiology.* 2010;6(3):325-42.

Blair SN; Wei M. Sedentary habits, health, and function in older women and men. *Am J Health Promot.* 2000;15(1):1-8.

Bowker SL; et al. Glucose-lowering agents and cancer mortality rates in type 2 diabetes: assessing effects of time-varying exposure. *Diabetologia.* 2010;53(8):1631-7.

Carlson SA; et al. Trend and prevalence estimates based on the 2008 Physical Activity Guidelines for Americans. *Am J Prev Med.* 2010;39(4):305-13.

Carnethon MR; et al. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *JAMA.* 2003;290(23):3092-100.

De Moor MH; et al. Genome-wide linkage scan for exercise participation in Dutch sibling pairs. *Eur J Hum Genet.* 2007;15(12):1252-9.

Enright PL; Sherrill DL. Reference equations for the six-minute walk test in healthy adults. *Am J Respir Crit Care Med.* 1998;158(5 Pt 1):1384-7.

Erickson KI; et al. Physical activity predicts gray matter volume in late adulthood: the cardiovascular health study. *Neurology.* 2010;75(16):1415-22.

Felitti VJ; et al. Obesity: problem, solution, or both? *Perm J.* 2010;14(1):24-30.

Ferney SL; et al. Awareness of and changing perceptions of physical activity guidelines among delegates at the Australian Conference of Science and Medicine in Sport. *J Sci Med Sport.* 2009;12(6):642-6.

Geldman M; Moore A; Cheek L. The effect of pre-injury physical fitness on the initial severity and recovery from whiplash injury, at six-month follow-up. *Clin Rehabil.* 2008;22(4):364-76.

Hanney WJ; et al. The influence of education and exercise on neck pain. *Am J Lifestyle Med.* 2010;4(2):166-75.

Katzmarzyk PT; Janssen I. The economic costs associated with physical inactivity and obesity in Canada: an update. *Can J Appl Physiol.* 2004;29(1):90-115.

Kodama S; et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA.* 2009;301(19):2024-35.

Landman GW; et al. Increased cancer mortality in type 2 diabetes. *Anticancer Res.* 2008;28(2B):1373-5.

Laukkanen JA; et al. Cardiorespiratory fitness, lifestyle factors and cancer risk and mortality in Finnish men. *Eur J Cancer.* 2010;46(2):355-63.

Lobelo F; Duperly J; Frank E. Physical activity habits of doctors and medical students influence their counseling practices. *Br J Sports Med.* 2009;43(2):89-92.

Loeppke R; et al. Health and productivity as a business strategy: a multiemployer study. *J Occup Environ Med.* 2009;51(4):411-28.

McAuley PA; et al. Obesity paradox and cardiorespiratory fitness in 12,417 male veterans aged 40 to 70 years. *Mayo Clin Proc.* 2010;85(2):115-21.

Phillips AC; Der G; Carroll D. Self-reported health, self-reported fitness, and all-cause mortality: prospective cohort study. *Br J Health Psychol.* 2010;15(Pt 2):337-46.

Sculco AD; et al. Effects of aerobic exercise on low back pain patients in treatment. *Spine J.* 2001;1(2):95-101.

Sigal RJ; et al. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med.* 2007;147(6):357-69.

Sinclair J; Lawson B; Burge F. Which patients receive advice on diet and exercise? Do certain characteristics affect whether they receive such advice? *Can Fam Physician.* 2008;54(3):404-12.

Stensel D. Primary prevention of CVD: physical activity. *Am Fam Phys.* 2010;82(2):136-38.

Storheim K; et al. Predictors of return to work in patients sick listed for sub-acute low back pain: a 12-month follow-up study. *J Rehabil Med.* 2005;37(6):365-71.

Teucher B; Rohrmann S; Kaaks R. Obesity: focus on all-cause mortality and cancer. *Maturitas*. 2010;65(2):112-6.

Van Strien T; Koenders P. How do physical activity, sports, and dietary restraint relate to overweight-associated absenteeism? *J Occup Environ Med*. 2010;52(9):858-64.

Warburton DE. Evidence-informed physical activity guidelines for Canadian adults. *Can J Public Health*. 2007;98(Suppl 2):S16-68.

Williams RM; et al. Effectiveness of workplace rehabilitation interventions in the treatment of work-related low back pain: a systematic review. *Disabil Rehabil*. 2007;29(8):607-24.

Ylipaa V; Arnetz BB; Preber H. Predictors of good general health, well-being, and musculoskeletal disorders in Swedish dental hygienists. *Acta Odont Scand*. 1999;57(5):277-82.